


Developing transgenic wheat and barley that exhibit resistance to *Fusarium* *graminearum* via glucoside conjugation of trichothecene mycotoxins

Gary J. Muehlbauer
University of Minnesota



Trichothecenes are virulence factors on wheat

Deoxynivalenol (DON)




Tri5 mutant – knockout in trichodiene synthase resulting in trichothecene nonproducing Fusarium



Proctor et al., 1995



UDP glucosyltransferase activity converts DON to DON-3-glucoside




HvUGT13248 encodes a barley UDP glucosyltransferase

- Confers resistance in yeast and Arabidopsis to DON
- Catalyzes the conjugation of DON to D3G
- Schweiger et al., 2010; Shin et al., 2012


Wheat transformation




Clemente collaboration




Transgenic wheat carrying *HvUGT13248* confer type I resistance




DON and D3G concentration changes in
HvUGT13248-#19 and BW from 1 to 21 dai



HvUGT13248
rapidly catalyzes
DON to D3G *in planta*






F. Berthiller collaboration




Transgenic wheat carrying *HvUGT1 3248* provide resistance to DON



Transgenic wheat carrying *HvUGT1 3248* provide resistance to DON




FHB disease phenotypes in field grown transgenic wheat (2013)




R. Dill-Macky collaboration

DON and D3G concentration in field grown transgenic wheat (2013)




Dill-Macky, Dong and Berthiller collaboration

Transgenic wheat carrying *HvUGT13248* confer FHB resistance (2014)




Dill-Macky and Dong collaboration


Introgression of HvUGT13248 into Rollag




UGT x Rollag BC₁F₂ lines point inoculation in the greenhouse




UGT x Rollag BC₁F₄ lines spray inoculated in the field





Transgenic wheat carrying *HvUGT13248* exhibit type II resistance to a NIV-producing *Fg* strain







UDP glucosyltransferase activity converts DON to DON-3-glucoside



HvUGT13248 detoxifies DON and NIV




Transgenic barley overexpressing *HvUGT13248* exhibits resistance to DON



Jochen Kumlehn collaboration

Transgenic barley overexpressing *HvUGT13248* exhibit resistance to DON



Summary

- We developed transgenic wheat lines carrying the *HvUGT13248* gene
- Transgenic wheat exhibited FHB resistance
- Transgenic wheat carrying *HvUGT13248* rapidly converted DON to D3G
- The *HvUGT13248* transgene in combination with *Fhb1* provides a reduction in severity, VSK, incidence and DON accumulation
- Transgenic wheat carrying *HvUGT13248* exhibited type II resistance to a NIV-*Fg* strain
- Transgenic barley carrying *HvUGT13248* exhibits resistance to DON



Acknowledgements

Muehlbauer lab

Jayanand Boddu

Seungho Cho

Stephanie Gardiner

Shane Heinen

Xin Li

Sanghyun Shin

USDA-ARS, St. Paul

Corby Kistler

University of Nebraska

Tom Clemente

IPK, Gatersleben

Jochum Kumlein



United States
Department of
Agriculture

National Institute
of Food and
Agriculture

U. of Minnesota

Jim Anderson

Ruth Dill-Macky

Yanhong Dong

USDA-ARS, Peoria, IL

Susan McCormick

Vienna U. of Technology

Christian Hametner

USDA-ARS, Fargo

Shaioman Chao

**University of Natural Resources and Life
Sciences, Austria**

Gerhard Adam

Wolfgang Schweiger

Marc Lemmens

Paula Kovalsky Paris

Juan Antonio Torres-Acosta

Franz Berthiller



Toxin concentrations in transgenic wheat carrying *HvUGT1 32248* in field trials. 2012-2014.

